

NAME OF THE MEDICINAL PRODUCT

Glucose 50 % B. Braun Concentrate for Solution for Infusion

COMPOSITION

100 ml of concentrate for formulation of solution for infusion contain

Glucose monohydrate	55.0 g
(equivalent to anhydrous glucose)	(50.0 g)

Excipient: Water for injection.

THERAPEUTIC INDICATIONS

Conditions caused by low glucose levels (hypoglycemia).

CONTRAINDICATIONS

Hyperglycemia, hypokalemia, acidosis.

UNDESIRABLE EFFECTS

Undesirable effects are listed according to their frequencies as follows:

Not known: (cannot be estimated from the available data)

Metabolism and nutrition disorders

Not known: Hypertonic dehydration, hospital-acquired hyponatremia*

Nervous system disorders

Not known: Hyponatremic encephalopathy*

General disorders and administration site conditions

Irritation at the administration site, vein irritation, thrombophlebitis, extravasation, local pain

*Hospital-acquired hyponatremia may cause irreversible brain injury and death due to the development of acute hyponatremic encephalopathy.

WARNINGS

Keep out of the sight and reach of children. Sterile and free of bacterial endotoxins. For single use only. Discard unused contents. Use only if concentrate is clear and container and closure are undamaged.

NOTE

Not all products are registered and approved for sale in all countries or regions. Indications of use may also vary by country and region. Please contact your country representative for product availability and information.

B. Braun Melsungen AG, 34212 Melsungen, Germany, 01/2018