

NAME OF THE MEDICINAL PRODUCT

Potassium Chloride 14.9 % Braun concentrate for solution for infusion

COMPOSITION

1 ml of concentrate for solution for infusion contains 0.149 g potassium chloride corresponding to 2 mmol K⁺ and 2 mmol Cl⁻

1 ampoule of 10 ml contains 1.49 g potassium chloride corresponding to 20 mmol K⁺ and 20 mmol Cl⁻

Excipient: Water for injection.

THERAPEUTIC INDICATIONS

Potassium deficiency, in particular when accompanied by hypochloreaemic alkalosis.

CONTRAINDICATIONS

Hyperkalemia; hyperchloremia.

Disorders frequently associated with hyperkalaemia, such as dehydration, impaired excretory renal function, Addison's disease, adynamia episodica hereditaria (Gamstorp disease), sickle cell anaemia.

UNDESIRABLE EFFECTS

Side effects are only to be expected as a result of absolute or relative overdose and/or excessively rapid administration rates. They correspond to the symptoms of hyperkalaemia. The frequency thereof is dose-dependent.

Metabolism and nutrition disorders

Acidosis, hyperkalaemia, hyperchloreaemia.

Cardiac disorders

Excessively rapid administration rates can lead to cardiac arrhythmias.

Gastro-intestinal disorders

Nausea.

WARNINGS

Keep out of the sight and reach of children. Only to be used diluted, as additive to infusion solutions. Discard unused contents.

NOTE

Prescription only

Not all products are registered and approved for sale in all countries or regions. Indications of use may also vary by country and region. Please contact your country representative for product availability and information.

B. Braun Melsungen AG, 34212 Melsungen, Germany, 07/2014