

NAME OF THE MEDICINAL PRODUCT

Sodium Bicarbonate 8.4% B.Braun solution for Infusion

COMPOSITION

1000 ml of solution for infusion contain:

Sodium hydrogen carbonate (sodium bicarbonate) 84.0 g

<i>Electrolyte concentrations</i>	<i>mmol/l</i>
Sodium	1000
Hydrogen carbonate	1000

Excipients: Edetate disodium (Ph. Eur.) 50 mg, water for injection.

THERAPEUTIC INDICATIONS

Correction of metabolic acidosis.

Urine alkalisation: In the case of intoxication with weak organic acids (e.g. barbiturates or acetylsalicylic acid); in order to improve the solubility of drug substances that are poorly soluble in neutral or acid medium (e.g. methotrexate, sulphonamides); in the case of haemolysis.

CONTRAINDICATIONS

Alkalosis, hypokalemia, hypernatremia.

UNDESIRABLE EFFECTS

The possible undesirable effects of Sodium Bicarbonate 8.4% B.Braun solution for Infusion result from the pharmacological and physical and chemical properties of the solution.

Undesirable effects are listed according to their frequency terms used in this section:

Not known: (cannot be estimated from the available data)

Metabolism and nutrition disorders

Not known: Administration can lead to hypernatremia and hyperosmolarity.

WARNINGS

Keep out of the sight and reach of children. For single use only. Discard container and unused content after use. Use only if solution is clear and colourless and container and closure are undamaged.

NOTE

Not all products are registered and approved for sale in all countries or regions. Indications of use may also vary by country and region. Please contact your country representative for product information and availability.

B. Braun Melsungen AG, 34212 Melsungen, Germany, 07/2014